


responding hydroxamic acids, fractionating these, and reconverting the pure oleohydroxamic acid to oleic acid.

The present procedure⁹ is a simplification of the methods of Brown and Shinowara⁵ and of Wheeler and Riemenschneider.⁶ It gives higher yields of a somewhat lower-quality product.

The procedure described is more satisfactory than those requiring the crystallization of soaps of oleic acid. For a review of these methods see ref. 5.

METHYL RICINOLEATE




Mol. wt. 312.5 (C₁₉H₃₆O₃)

Submitted by DANIEL SWERN and E. F. JORDAN, JR., Eastern Regional Research Laboratory, Philadelphia.

Checked by L. H. MASON and H. E. CARTER, University of Illinois, Urbana.

I. Principle



II. Procedure

One thousand grams of U.S.P. castor oil are added to a solution of 3 gm. of metallic sodium in 2 l. of anhydrous methanol in a 12-l. Pyrex round-bottomed flask. The reaction mixture is refluxed for 45 minutes and then diluted, with stirring, with about 7 l. of water containing 25 ml. of 6 *N* hydrochloric acid. The lower aqueous layer is siphoned off and discarded. The methyl esters are transferred to a 4-l. separatory funnel and washed with water until free of chloride ion. The crude, washed esters

⁹ D. Swern, H. B. Knight, and T. W. Findley, *Oil & Soap*, **21**, 133 (1944).

are dried by heating to 100–105° under moderate vacuum in a stream of inert gas. The yield at this point is 970–980 gm. of a pale yellow oil. This is fractionally distilled *in vacuo*.¹ The fraction distilling at 185–187°/2 mm. consists of essentially pure methyl ricinoleate. The yield is 670–680 gm. (84–85% of the theory based on the assumption that castor oil contains 80% of ricinoleic acid).

III. Properties and Purity of Product

Methyl ricinoleate is a colorless, odorless oil melting at –6°. It gives the following constants: refractive index, $n_D^{20} = 1.463$; specific rotation, $[\alpha]_D^{25} = +5.05^\circ$ (5% solution in acetone); iodine number, 81–82; saponification equivalent, 312–313; hydroxyl, 5.5%.

IV. Methods of Preparation

Methyl ricinoleate was prepared by methanolysis of castor oil followed by fractional distillation and low-temperature solvent crystallization.² The present procedure is that of Kass and Radlove.³

¹ A column with a low pressure-drop should be used to avoid pot temperatures much in excess of 220°. In addition, the column should have a fairly high throughput to keep the distillation time as short as possible. A Vigreux column, 3 ft. long and 1 in. in diameter (five theoretical plates), electrically heated and well insulated, is satisfactory. The checkers found that a vacuum-jacketed silvered Vigreux column gave lower yields than an electrically heated one.

² J. B. Brown and N. D. Green, *J. Am. Chem. Soc.*, **62**, 738 (1940).

³ J. P. Kass and S. B. Radlove, *J. Am. Chem. Soc.*, **64**, 2253 (1942).